

## CULTURE ADVICE

### pac® Duftpelargonien® [scented-leaved pelargoniums]

<b>Description:</b>	It is primarily the leaves of the scented-leaved pelargoniums that are of importance; a safe flowering at the main selling time - weeks 17 to 19 - can only be expected with the P.fragrans, Concolor Lace and Attar of Roses varieties; all other varieties usually start blooming at a later time. If young plants are purchased early enough - between calendar weeks 48 and 2 - the producer may insert a cooling period after potting which improves flower formation. Scented-leaved pelargoniums should as a rule be sold at a higher price than P. zonals and P. peltatum varieties, and they should be given a preferred place in the selling area.
<b>Potting:</b>	Young plants are potted in 13-15 cm pots (if they are purchased as from week 7 or 8).
<b>Substrate:</b>	Loose, well-drained substrate (with increased white peat portion) or standard soil of type 2 (type T). Own mixtures with a compost portion may also be used. In any case, the soil shall be coarse, permeable and free from diseases. N: 150 - 250 mg/l P: 130 - 200 mg/l K: 300 - 350 mg/l pH: 5.8 - 6.0 EC: 1.0 - 1.2 mS
<b>Fertilization:</b>	Liquid fertilization at regular intervals: 1 g/l, or 1 to 2 times a week: max. 2.5 g/l
<b>Temperature:</b>	Cool cultivation at 12 °C is energy-saving.
<b>Flowering:</b>	As from mid-May.
<b>Growth regulators:</b>	The following should be observed during cultivation: Move the plants in due time, do not set them too close, apply CCC at regular intervals. Basically, our products can be subdivided into two growth types: a) strong growth (0.15 % per week) Sweet Mimosa, Clorinda, Attar of Roses, Pink Capitatum, P.citriodorum, P.quercifolium, Torento b) weak and compact growth (treat as required) Orange Fizz, Lady Plymouth, Concolor Lace, P.x fragrans
<b>Pinching:</b>	With the exception of P.x Fragrans, the plants need to be pinched soon to get a good quality. A pinching of the other species/varieties is not absolutely necessary. However, if branching is not satisfactory, some trimming or shaping is recommended to get a better plant build-up. 1 to 2 weeks after potting or immediately in case of a suitable young plant size.
<b>Diseases / pest:</b>	Botrytis primarily with Concolor Lace, white fly primarily with Orange Fizz

Water balance is of importance; the plants must not stand too wet, danger of botrytis, damage to roots and irregular plant build-up are the consequences.

Updated: 08/09/2009